



Osmosis


By using this tool, you will be able to visualise the process of osmosis which gives a clear understating of this process.


Please go to the URL
<https://pbslm-contrib.s3.amazonaws.com/WGBH/arct15/SimBucket/Simulations/osmosis/content/index.html>
(or)
Scan the QR code on the right side



Steps

- Open the Browser and type the URL given (or) Scan the QR Code.
- Now you will see the webpage as shown below.
- The two compartments were separated by a semipermeable membrane.
- On the left side (Box 2) contains pure solvent (water) and right side contains solution (solute + water).
- The number of solvent molecules (water) on each side is shown in the area designated by box 3 and 5.
- Now you can see the water molecules moving from solvent side to solution side through the membrane. It reflects in the change in number of water molecules on either side.
- You can restart the simulation by clicking the reset button in box 1





Linus Carl Pauling was an American chemist, biochemist, peace activist, author and educator. In addition to his contribution to chemistry and he also worked with many biologists.

He received the Nobel Prize in Chemistry in 1954 for his research into the nature of the chemical bond and its application to the elucidation of the structure of complex substances.

Learning Objectives

After studying this unit students will be able to

- describe Kossel – Lewis approach to chemical bonding
- explain the octet rule
- sketch the Lewis structures of simple molecules
- describe the formation of different types of bonds and bond parameters
- sketch the resonance structures for simple molecules
- apply the concept of electronegativity to explain the polarity of covalent bonds
- describe VSEPR theory and predict the shapes of simple molecules
- explain the valence bond approach for the formation of covalent bonds
- explain the different types of hybridisation involving s, p & d orbitals and sketch shapes of simple covalent molecules
- explain the molecular orbital theory, calculate the bond order and explain the magnetic properties of H_2 , O_2 , N_2 , CO and NO
- describe metallic bonding briefly.

10.1 Introduction

Diamond is very hard while its allotrope graphite is very soft. Gases like hydrogen and oxygen are diatomic while the inert gases are monoatomic.


Carbon combines with chlorine to form carbon tetrachloride, which is a liquid and insoluble (immiscible) in water. Sodium combines with chlorine atom to form sodium chloride, a hard and brittle compound that readily dissolves in water. The possible reason for these observations lies in the type of interaction that exists between the atoms of these molecules and these interactions are responsible for holding the atoms/ions together. The interatomic attractive forces which hold the constituent atoms/ions together in a molecule are called chemical bonds.

Why do atoms combine only in certain combinations to form molecules? For example oxygen combines with hydrogen to give water (H₂O) and with carbon it gives carbon dioxide (CO₂). The structure of water is 'V' shaped while that of the carbon dioxide is linear. Such questions can be answered using the principles of chemical bonding. In this unit we will analyse the various theories and their principles, which were developed over the years to explain the nature of chemical bonding.

10.1.1 Kossel – Lewis approach to chemical bonding

A logical explanation for chemical bonding was provided by Kossel and Lewis in 1916. Their approach to chemical bonding is based on the inertness of the noble gases which have little or no tendency to combine with other atoms. They proposed that the noble gases are stable due to their completely filled outer shell electronic configuration. Elements other than noble gases, try to attain the completely filled electronic configurations by losing, gaining

or sharing one or more electrons from their outer shell. For example, sodium loses one electron to form Na⁺ ion and chlorine accepts that electron to give chloride ion (Cl⁻), enabling both atoms to attain the nearest noble gas configuration. The resultant ions, Na⁺ and Cl⁻ are held together by electrostatic attractive forces and the attractive force is called a chemical bond, more specifically an electrovalent bond.



G. N. Lewis proposed that the attainment of stable electronic configuration in molecules such as diatomic nitrogen, oxygen etc... is achieved by mutual sharing of the electrons. He introduced a simple scheme to represent the chemical bond and the electrons present in the outer shell of an atom, called Lewis dot structure. In this scheme, the valence electrons (outer shell electrons) of an element are represented as small dots around the symbol of the element. The first four valence electrons are denoted as single dots around the four sides of the atomic symbol and then the fifth onwards, the electrons are denoted as pairs. For example, the electronic configuration of nitrogen is 1s², 2s², 2p³. It has 5 electrons in its outer shell (valence shell). The Lewis structure of nitrogen is as follows.



Fig 10.1 Lewis Structure of Nitrogen atom

Similarly, Lewis dot structure of carbon, oxygen can be drawn as shown below.



Fig 10.2 Lewis Structures of C & O atoms

Only exception to this is helium which has only two electrons in its valence shell which is represented as a pair of dots (duet).



Fig 10.3 Lewis Structures of He atom

Octet rule

The idea of Kossel – Lewis approach to chemical bond lead to the octet rule, which states that “*the atoms transfer or share electrons so that all atoms involved in chemical bonding obtain 8 electrons in their outer shell (valence shell)*”.

10.2 Types of chemical bonds

The chemical bonds can be classified based on the nature of the interaction between the bonded atoms. Two major types of chemical bonds are covalent bonds and ionic bonds. Generally metals reacts with non-metals to form ionic compounds, and the covalent bonds are present in the compounds formed by nonmetals.

10.2.1 Covalent bonds:

Do you know all elements (except noble gases) occurs either as compounds or as polyatomic molecules? Let us consider hydrogen gas in which two hydrogen atoms bind to give a dihydrogen molecule. Each hydrogen atom has one electron and it requires one more electron to attain the electronic configuration of the nearest noble gas helium. Lewis suggested that both hydrogen atoms will attain the stable

configuration by mutually sharing the electrons available with them. Similarly, in the case of oxygen molecule, both the oxygen atoms share two electron pairs between them and in nitrogen molecule three electron pairs are shared between two nitrogen atoms. *This type of mutual sharing of one or more pairs of electrons between two combining atoms results in the formation of a chemical bond called a covalent bond.* If two atoms share just one pair of electron a single covalent bond is formed as in the case of hydrogen molecule. If two or three electron pairs are shared between the two combining atoms, then the covalent bond is called a double bond or a triple bond, respectively.




Fig 10.4 Representation of Lewis Structures of covalent bonds

10.2.2 Representing a covalent bond - Lewis structure (Lewis dot structure)

Lewis structure (Lewis dot structure) is a pictorial representation of covalent bonding between the combining atoms. In this structure the shared valence electrons are represented as a pair of dots between the combining atoms and the unshared electrons of the atoms are represented as a pair of dots (lone pair) on the respective individual atoms.

The Lewis dot structure for a given compound can be written by following the steps given below. Let us understand these steps by writing the Lewis structure for water.

1. **Draw the skeletal structure of the molecule.** In general, the less electronegative atom is placed at the centre. Hydrogen and fluorine atoms should be placed at the terminal positions. For water, the skeletal structure is



2. **Calculate the total number of valence electrons of all the atoms in the molecule.** In case of polyatomic ions the charge on ion should also be considered during the calculation of the total number of valence electrons. In case of anions the number of negative charges should be added to the number of valence electrons. For positive ions the total number of positive charges should be subtracted from the total number of valence electrons.

In water, total number of valence electron = $[2 \times 1$ (valence electron of

hydrogen)] + $[1 \times 6$ (valence electrons of oxygen)] = $2 + 6 = 8$.

3. **Draw a single bond between the atoms in the skeletal structure of the molecule.** Each bond will account for two valence electrons (a bond pair). For water, we can draw two bonds accounting for four valence electrons as follows.



4. **Distribute the remaining valence electrons as pairs (lone pair), giving octet (only duet for hydrogen) to the atoms in the molecule.** The distribution of lone pairs starts with the most electronegative atoms followed by other atoms.

In case of water, the remaining four electrons (two lone pairs) are placed on the most electronegative central oxygen, giving octet.



5. **Verify whether all the atoms satisfy the octet rule (for hydrogen duet).** If not, use the lone pairs of electrons to form additional bond to satisfy the octet rule.

In case of water, oxygen has octet and the hydrogens have duets, hence there is no need for shifting the lone pairs. The Lewis structure of water is as follows



Fig 10.5 Lewis structure of water

Let us draw the Lewis structure for nitric acid.

1. Skeletal structure




- Total number of valence electrons in HNO_3
 $= [1 \times 1(\text{hydrogen})] + [1 \times 5(\text{nitrogen})] + [3 \times 6(\text{oxygen})] = 1 + 5 + 18 = 24$
- Draw single bonds between atoms. Four bonds can be drawn as shown in the figure for HNO_3 which account for eight electrons (4 bond pairs).



- Distribute the remaining sixteen ($24 - 8 = 16$) electrons as eight lone pairs starting from most electronegative atom, the oxygen. Six lone pairs are distributed to the two terminal oxygens

(three each) to satisfy their octet and two pairs are distributed to the oxygen that is connected to hydrogen to satisfy its octet.



- Verify whether all the atoms have octet configuration. In the above distribution, the nitrogen has one pair short for octet. Therefore, move one of the lone pair from the terminal oxygen to form another bond with nitrogen.

The Lewis structure of nitric acid is given as




Fig 10. 6 Lewis structure of Nitric acid

Table 10.1 : The Lewis dot structures for some molecules

S. No	Molecule	Lewis Structure	
1.	Sulphur trioxide (SO_3)	$\begin{array}{c} \ddot{\text{O}} \\ \\ \ddot{\text{O}} - \text{S} = \ddot{\text{O}} \end{array}$	$\begin{array}{c} \ddot{\text{O}} \\ \ddot{\text{O}} : \text{S} : \ddot{\text{O}} \end{array}$
2.	Ammonia (NH_3)	$\begin{array}{c} \text{H} \\ \\ \text{H} - \text{N} - \text{H} \\ \vdots \end{array}$	$\begin{array}{c} \text{H} \\ \text{H} : \text{N} : \text{H} \\ \vdots \end{array}$
3.	Methane (CH_4)	$\begin{array}{c} \text{H} \\ \\ \text{H} - \text{C} - \text{H} \\ \\ \text{H} \end{array}$	$\begin{array}{c} \text{H} \\ \text{H} : \text{C} : \text{H} \\ \vdots \\ \text{H} \end{array}$
4.	Dinitrogen Pentoxide (N_2O_5)	$\begin{array}{c} \ddot{\text{O}} = \text{N} - \ddot{\text{O}} - \text{N} = \ddot{\text{O}} \\ \quad \quad \\ \ddot{\text{O}} \quad \quad \ddot{\text{O}} \end{array}$	$\begin{array}{c} \ddot{\text{O}} : \text{N} : \ddot{\text{O}} : \text{N} : \ddot{\text{O}} \\ \ddot{\text{O}} : \quad \ddot{\text{O}} : \end{array}$

Note

It is to be noted that nearly in all their compounds, certain elements form a fixed number of bonds. For example, Fluorine forms only one bond. Hydrogen, oxygen, nitrogen and carbon atoms form one, two, three and four bonds, respectively.

Evaluate Yourself

- 1) Draw the Lewis structures for
 - i) Nitrous acid (HNO_2)
 - ii) Phosphoric acid
 - iii) Sulphur trioxide (SO_3)

10.2.3 Formal charge:

Let us draw the Lewis structure for carbon dioxide.

1. Skeletal structure



2. Total number of valence electrons in CO_2

$$= [1 \times 4(\text{carbon})] + [2 \times 6(\text{oxygen})] = 4 + 12 = 16$$

3. Draw single bonds between atoms. Two bonds can be drawn as shown in the figure for CO_2 which accounts for four electrons (2 bond pairs).



4. Distribute the remaining twelve electrons ($16 - 4 = 12$) as six lone pairs starting from most electronegative atom, the oxygen. Six lone pairs are distributed to the two terminal oxygens (three each) to satisfy their octet.



5. Verify whether all the atoms have octet configuration. In the above distribution, the central carbon has two pair short

for octet. Therefore, to satisfy the octet rule two lone pairs from one oxygen or one pair from each oxygen can be moved to form multiple bonds, leading the formation of two possible structures for carbon dioxide as shown below




Fig 10.7 (a) two possible structures for carbon dioxide

Similarly, the Lewis structure for many molecules drawn using the above steps gives more than one acceptable structure. Let us consider the above mentioned two structures of carbon dioxide.

Which one the above forms represents the best distribution of electrons in the molecule. To find an answer, we need to know the formal charge of each atom in the Lewis structures. Formal charge of an atom in a molecule, is the electrical charge difference between the valence electron in an isolated atom and the number of electrons assigned to that atom in the Lewis structure.

$$\text{Formal charge of an atom} = N_v - \left(N_l + \frac{N_b}{2} \right)$$

Where,

N_v - Number of valence electron of atom in its isolated state.

N_l - Number of electrons present as lone pairs around the atom in the Lewis structure

N_b - Number of electrons present in bonds around the atom (bond pairs) in the Lewis structure]

Now let us calculate the formal charge on all atoms in both structures,

For Structure 1,

$$\begin{aligned}\text{Formal charge on carbon} &= N_v - \left(N_l + \frac{N_b}{2} \right) \\ &= 4 - \left(0 + \frac{8}{2} \right) = 0\end{aligned}$$

$$\begin{aligned}\text{Formal charge on oxygen} &= 6 - \left(4 + \frac{4}{2} \right) \\ &= 0 \text{ (for both oxygens)}\end{aligned}$$

For structure 2

Formal charge on carbon

$$\begin{aligned}&= N_v - \left(N_l + \frac{N_b}{2} \right) \\ &= 4 - \left(0 + \frac{8}{2} \right) = 0\end{aligned}$$

Formal charge on singly bonded oxygen

$$= 6 - \left(6 + \frac{2}{2} \right) = -1$$

Formal charge on triply bonded oxygen

$$= 6 - \left(2 + \frac{6}{2} \right) = +1$$




Fig 10. 7 (b) two possible structures for carbon dioxide (with formal charges)

After calculating the formal charges, the best representation of Lewis structure can be selected by using following guidelines.

1. A structure in which all formal charges are zero preferred over the one with charges.
2. A structure with small formal charges

is preferred over the one with higher formal charges.

3. A structure in which negative formal charges are placed on the most electronegative atom is preferred.

In case of CO₂ structures, the structure one is preferred over the structure 2 as it has zero formal charges for all atoms.

10.2.4 Lewis structures for exceptions to octet rule

The octet rule is useful for writing Lewis structures for molecules with second period element as central atoms. In some molecules, the central atoms have less than eight electrons around them while some others have more than eight electrons. Exception to the octet rule can be categorized into following three types.

1. Molecules with electron deficient central atoms
2. Molecules containing odd electrons
3. Molecules with expanded valence shells

Molecules with electron deficient central atoms

Let us consider boron trifluoride, as an example. The central atom boron has three valence electron and each fluorine has seven valence electrons. The Lewis structure is




Fig 10. 8 (a) Lewis structure of BF₃

In the above structure, only six electrons around boron atom. Moving a

lone pair from one of the fluorine to form additional bond as shown below.




Fig 10. 8 (b) Lewis structure of BF_3

However, the above structure is unfavourable as the most electronegative atom fluorine shows positive formal charge and hence the structure with incomplete octet is the favourable one. Molecules such as BCl_3 , BeCl_2 , etc... also have incomplete octets.

Molecules containing odd electrons

Few molecules have a central atom with an odd number of valence electrons. For example, in nitrogen dioxide and nitric oxide all the atoms does not have octet configuration. The lewis structure of the above molecules are shown in the figure.




Fig 10. 9 Lewis structures of Nitric oxide and Nitrogen dioxide (with formal charges)

Molecules with expanded valence shells

In molecules such as sulphur hexafluoride (SF_6), phosphorous pentachloride (PCl_5) the central atom has more than eight valence electrons around them. Here the central atom can accommodate additional electron pairs by using outer vacant d orbitals. In SF_6 the central atom sulphur is surrounded by six bonding pair of electrons or twelve electrons.




Fig 10. 10 Lewis structures for SF_6 and PCl_5


Evaluate Yourself

- Calculate the formal charge on each atom of carbonyl chloride (COCl_2)

10.3 Ionic or electrovalent bond

When the electronegativity difference between the two combining atoms is large, the least electronegative atom completely transfers one or more of its valence electrons to the other combining atom so that both atoms can attain the nearest inert gas electronic configuration. The complete transfer of electron leads to the formation of a cation and an anion. Both these ions are held together by the electrostatic attractive force which is known as ionic bond.

Let us consider the formation potassium chloride. The electronic configuration of potassium and chlorine are



Chlorine (Cl) : [Ne]3s², 3p⁵

Potassium has one electron in its valence shell and chlorine has seven electron in its valence shell. By losing one electron potassium attains the inert gas electronic configuration of argon and becomes a unipositive cation (K⁺) and chlorine accepts this electron to become uninegative chloride ion (Cl⁻) there by attaining the stable electronic configuration of argon. These two ions combine to form an ionic crystal in which they are held together by electrostatic attractive force. The energy required for the formation of one mole of K⁺ is 418.81 kJ (ionization energy) and the energy released during the formation of one mole of Cl⁻ is -348.56 kJ (electron gain enthalpy). The sum of these two energies is positive (70.25 kJ) However, during the formation of one mole potassium chloride crystal from its constituent ions, 718 kJ energy is released. This favours the formation of KCl and its stability.

Evaluate Yourself

- 3) Explain the ionic bond formation in MgO and CaF₂

10.4 Coordinate covalent bond

In the formation of a covalent bond, both the combining atoms contribute one electron each and these electrons are mutually shared among them. However, in certain bond formation, one of the combining atoms donates a pair of electrons i.e. two electrons which are necessary for the covalent bond formation, and these electrons are shared by both the combining atoms. These type of bonds are called coordinate

covalent bond or coordinate bond. The combining atom which donates the pair of electron is called a donor atom and the other atom an acceptor atom. This bond is denoted by an arrow starting from the donor atom pointing towards the acceptor atom. (Later in coordination compound, we will refer the donor atom as ligand and the acceptor atom as central-metal atom/ion.

For Example, in ferrocyanide ion [Fe(CN)₆]⁴⁻, each cyanide ion (CN⁻) donates a pair of electrons to form a coordinate bond with iron (Fe²⁺) and these electrons are shared by Fe²⁺ and CN⁻.




Fig 10.11 Structure of Ferrocyanide ion

In certain cases, molecules having a lone pair of electrons such as ammonia donates its pair to an electron deficient molecules such as BF₃. to form a coordinate




Fig 10.12 Structure of BF₃ → NH₃

10.5 Bond parameters

A covalent bond is characterised by parameters such as bond length, bond angle, bond order etc... A brief description of some of the bond parameters is given below.

10.5.1 Bond length

The distance between the nuclei of the two covalently bonded atoms is called bond length. Consider a covalent molecule A-B. The bond length is given by the sum of the radii of the bonded atoms ($r_A + r_B$). The length of a bond can be determined by spectroscopic, x-ray diffraction and electron-diffraction techniques. The bond length depends on the size of the atom and the number of bonds (multiplicity) between the combining atoms.




Fig 10.13 Bond length of covalent molecule A-B

Greater the size of the atom, greater will be the bond length. For example, carbon-carbon single bond length (1.54 \AA) is longer than the carbon-nitrogen single bond length (1.43 \AA).

Increase in the number of bonds between the two atoms decreases the bond

length. For example, the carbon-carbon single bond is longer than the carbon-carbon double bond (1.33 \AA) and the carbon-carbon triple bond (1.20 \AA).

10.5.2 Bond order

The number of bonds formed between the two bonded atoms in a molecule is called the bond order. In Lewis theory, the bond order is equal to the number of shared pair of electrons between the two bonded atoms. For example in hydrogen molecules, there is only one shared pair of electrons and hence, the bond order is one. Similarly, in H_2O , HCl , Methane, etc the central atom forms single bonds with bond order of one.

Table 10.2 Bond order of some common bonds:

S. No.	Molecule	Bonded atoms	Bond order (No. of shared pair of electrons between bonded atoms)
1	H_2	H-H	1
2	O_2	O=O	2
3	N_2	$\text{N}\equiv\text{N}$	3
4	HCN	$\text{C}\equiv\text{N}$	3
5	HCHO	C=O	2
6	CH_4	C-H	1
7	C_2H_4	C=C	2

10.5.3 Bond angle

Covalent bonds are directional in nature and are oriented in specific directions in space. This directional nature creates a fixed angle between two covalent bonds in a molecule and this angle is termed as bond angle. It is usually expressed in degrees. The bond angle can be determined by spectroscopic methods and it can give some idea about the shape of the molecule.

Table 10.3 Bond angles for some common molecules



S. No.	Molecule	Atoms defining the angle	Bond angle ($^{\circ}$)
1	CH ₄	H-C-H	109 $^{\circ}$ 28'
2	NH ₃	H-N-H	107 $^{\circ}$ 18'
3	H ₂ O	H-O-H	104 $^{\circ}$ 35'

10.5.4 Bond enthalpy

The bond enthalpy is defined as the minimum amount of energy required to break one mole of a particular bond in molecules in their gaseous state. The unit of bond enthalpy is kJ mol⁻¹. Larger the bond enthalpy, stronger will be the bond. The bond energy value depends on the size of the atoms and the number of bonds between the bonded atoms. Larger the size of the atom involved in the bond, lesser is the bond enthalpy.

In case of polyatomic molecules with, two or more same bond types, in the term average bond enthalpy is used. For such

bonds, the arithmetic mean of the bond energy values of the same type of bonds is considered as average bond enthalpy. For example in water, there are two OH bonds present and the energy needed to break them are not same.



The average bond enthalpy of OH bond in water = $\frac{502+427}{2} = 464.5 \text{ kJ mol}^{-1}$


Table 10.4 Bond lengths and bond enthalpies of some common bonds:

S. No.	Bond type	Bond Enthalpy (kJ mol ⁻¹)	Bond Length (Å)
1	H-H	432	0.74
2	H-F	565	0.92
3	H-Cl	427	1.27
4	H-Br	363	1.41
5	H-I	295	1.61
6	C-H	413	1.09
7	C-C	347	1.54
8	C-Si	301	1.86
9	C-N	305	1.47
10	C-O	358	1.43
11	C-P	264	1.87
12	C-S	259	1.81
13	C-F	453	1.33
14	C-Cl	339	1.77
15	C-Br	276	1.94
16	C-I	216	2.13

10.5.5 Resonance


When we write Lewis structures for a molecule, more than one valid Lewis structures are possible in certain cases. For example let us consider the Lewis structure of carbonate ion $[\text{CO}_3]^{2-}$.

The skeletal structure of carbonate ion (The oxygen atoms are denoted as O_A , O_B & O_C



Total number of valence electrons = $[1 \times 4(\text{carbon})] + [3 \times 6(\text{oxygen})] + [2(\text{charge})]$
= 24 electrons.

Distribution of these valence electrons gives us the following structure.



Complete the octet for carbon by moving a lone pair from one of the oxygens (O_A) and write the charge of the ion (2-) on the upper right side as shown in the figure.




Fig 10.14 (a) Lewis Structure of CO_3^{2-}

In this case, we can draw two additional Lewis structures by moving the lone pairs from the other two oxygens

(O_B and O_C) thus creating three similar structures as shown below in which the relative position of the atoms are same. They only differ in the position of bonding and lone pair of electrons. Such structures are called resonance structures (canonical structures) and this phenomenon is called resonance.




Fig 10.14 (b) Resonance structures of CO_3^{2-}

It is evident from the experimental results that all carbon-oxygen bonds in carbonate ion are equivalent. The actual structure of the molecules is said to be the resonance hybrid, an average of these three resonance forms. It is important to note that carbonate ion does not change from one structure to another and vice versa. It is not possible to picturise the resonance hybrid by drawing a single Lewis structure. However, the following structure gives a qualitative idea about the correct structure.




Fig 10. 14 (c) Resonance Hybrid structures of CO_3^{2-}

It is found that the energy of the resonance hybrid (structure 4) is lower than that of all possible canonical structures (Structure 1, 2 & 3). The difference in energy between structure 1 or 2 or 3, (most stable canonical structure) and structure 4 (resonance hybrid) is called resonance energy.

Evaluate Yourself

4) Write the resonance structures for

- i) Ozone molecule ii) N_2O

10.5.6 Polarity of Bonds

Partial ionic character in covalent bond:

When a covalent bond is formed between two identical atoms (as in the case of H_2 , O_2 , Cl_2 etc...) both atoms have equal tendency to attract the shared pair of electrons and hence the shared pair of electrons lies exactly in the middle of the nuclei of two atoms. However, in the case of covalent bond formed between atoms having different electronegativities, the atom with higher electronegativity will have greater tendency to attract the shared pair of electrons more towards itself than the other atom. As a result the cloud of shared electron pair gets distorted.

Let us consider the covalent bond between hydrogen and fluorine in hydrogen

fluoride. The electronegativities of hydrogen and fluorine on Pauling's scale are 2.1 and 4 respectively. It means that fluorine attracts the shared pair of electrons approximately twice as much as the hydrogen which leads to partial negative charge on fluorine and partial positive charge on hydrogen. Hence, the H-F bond is said to be polar covalent bond.

Here, a very small, equal and opposite charges are separated by a small distance (91 pm) and is referred to as a dipole.

Dipole moment:

The polarity of a covalent bond can be measured in terms of dipole moment which is defined as

$$\mu = q \times 2d$$

Where μ is the dipole moment, q is the charge and $2d$ is the distance between the two charges. The dipole moment is a vector and the direction of the dipole moment vector points from the negative charge to positive charge.





Fig 10. 15 Representation of Dipole

The unit for dipole moment is coulomb meter (C m). It is usually expressed in Debye unit (D). The conversion factor is $1 \text{ Debye} = 3.336 \times 10^{-30} \text{ C m}$

Diatomic molecules such as H_2 , O_2 , F_2 etc... have zero dipole moment and are called non polar molecules and molecules such as HF, HCl, CO, NO etc... have non zero dipole moments and are called polar molecules.

Molecules having polar bonds will not necessarily have a dipole moment. For example, the linear form of carbon dioxide has zero dipole moment, even though it has two polar bonds. In CO_2 , the dipole moments of two polar bonds (CO) are equal in magnitude but have opposite direction. Hence, the net dipole moment of the CO_2 is,

$$\mu = \mu_1 + \mu_2 = \mu_1 + (-\mu_1) = 0$$


In case of water net dipole moment is the vector sum of $\mu_1 + \mu_2$ as shown.




Fig 10. 16 Dipole moment in water

Dipole moment in water is found to be 1.85D

Table 10. 5 Dipole moments of common molecules

S. No.	Molecule	Dipole moment (in D)
1	HF	1.91
2	HCl	1.03
3	H_2O	1.85
4	NH_3	1.47
5	CHCl_3	1.04

The extent of ionic character in a covalent bond can be related to the electronegativity difference to the bonded atoms. In a typical polar molecule, $\text{A}^{\delta-}\text{-B}^{\delta+}$, the electronegativity difference ($\chi_A - \chi_B$) can be used to predict the percentage of ionic character as follows.

If the electronegativity difference ($\chi_A - \chi_B$), is

equal to 1.7, then the bond A-B has 50% ionic character

if it is greater than 1.7, then the bond A-B has more than 50% ionic character,

and if it is lesser than 1.7, then the bond A-B has less than 50% ionic character.

Evaluate Yourself

- 5) Of the two molecules OCS and CS_2 which one has higher dipole moment value? why?

Partial covalent character in ionic bonds:

Like the partial ionic character in covalent compounds, ionic compounds show partial covalent character. For example, the ionic compound, lithium chloride shows covalent character and is soluble in organic solvents such as ethanol.

The partial covalent character in ionic compounds can be explained on the basis of a phenomenon called polarisation. We know that in an ionic compound, there is an electrostatic attractive force between the cation and anion. The positively charged cation attracts the valence electrons of anion while repelling the nucleus. This causes a distortion in the electron cloud of the anion and its electron density drifts towards the cation, which results in some sharing of the

valence electrons between these ions. Thus, a partial covalent character is developed between them. This phenomenon is called polarisation.

The ability of a cation to polarise an anion is called its polarising ability and the tendency of an anion to get polarised is called its polarisability. The extent of polarisation in an ionic compound is given by the Fajans rules

Fajans Rules

- (i) To show greater covalent character, both the cation and anion should have high charge on them. Higher the positive charge on the cation, greater will be the attraction on the electron cloud of the anion. Similarly higher the magnitude of negative charge on the anion, greater is its polarisability. Hence, the increase in charge on cation or in anion increases the covalent character

Let us consider three ionic compounds aluminum chloride, magnesium chloride and sodium chloride. Since the charge of the cation increase in the order $\text{Na}^+ < \text{Mg}^{2+} < \text{Al}^{3+}$, the covalent character also follows the same order $\text{NaCl} < \text{MgCl}_2 < \text{AlCl}_3$.

- (ii) The smaller cation and larger anion show greater covalent character due to the greater extent of polarisation.

Lithium chloride is more covalent than sodium chloride. The size of Li^+ is smaller than Na^+ and hence the polarising power of Li^+ is more. Lithium iodide is more covalent than lithium chloride as the size of I^- is larger than the Cl^- . Hence I^- will be more polarised than Cl^- by the cation, Li^+ .

- (iii) Cations having $ns^2 np^6 nd^{10}$ configuration show greater polarising power than the cations with $ns^2 np^6$ configuration. Hence, they show greater covalent character.

CuCl is more covalent than NaCl . Compared to Na^+ (1.13 Å). Cu^+ (0.6 Å) is small and have $3s^2 3p^6 3d^{10}$ configuration.

Electronic configuration of Cu^+
[Ar] $3d^{10}$

Electronic Configuration of Na^+
[He] $2s^2, 2p^6$


10.6 Valence Shell Electron Pair Repulsion (VSEPR) theory

Lewis concept of structure of molecules deals with the relative position of atoms in the molecules and sharing of electron pairs between them. However, we cannot predict the shape of the molecule using Lewis concept. Lewis theory in combination with VSEPR theory will be useful in predicting the shape of molecules.

Important principles of VSEPR Theory are as follows:

1. The shape of the molecules depends on the number of valence shell electron pair around the central atom.
2. There are two types of electron pairs namely bond pairs and lone pairs. The bond pair of electrons are those shared between two atoms, while the lone pairs are the valence electron pairs that are not involved in bonding.
3. Each pair of valence electrons around the central atom repels each other and hence, they are located as far away as possible in three dimensional space to minimize the repulsion between them.









4. The repulsive interaction between the different types of electron pairs is in the following order.























The lone pair of electrons are localised only on the central atom and interacts with only one nucleus whereas the bond pairs are shared between two atoms and they interact with two nuclei. Because of this the lone pairs occupy more space and have greater repulsive power than the bond pairs in a molecule.

The following Table illustrates the shapes of molecules predicted by VSEPR theory. Consider a molecule AB_x where A is the central atom and x represents the number of atoms of B covalently bonded to the central atom A. The lone pairs present in the atoms are denoted as L.

Table 10. 6 Shapes of molecules predicted by VSEPR theory.

Number of electron Pairs	Molecule	No. of bond pairs	No. of lone pairs	Shape	Molecular geometry	Examples
2	AB_2	2	-	 Linear	 Linear	$BeCl_2$, $HgCl_2$, CO_2 , CS_2 , HCN , BeF_2
3	AB_3	3	-	 Trigonal planar	 Trigonal planar	BF_3 , BCl_3 , NO_3^- , BF_3 , CO_3^{2-} , $HCHO$
	AB_2L	2	1	 Bent or V - Shape	 Bent or V - Shape	SO_2 , O_3 , $PbCl_2$, $SnBr_2$
4	AB_4	4	-	 Tetrahedral	 Tetrahedral	CH_4 , CCl_4 , CCl_2F_2 , SO_4^{2-} , ClO_4^- , NH_4^+

Number of electron Pairs	Molecule	No. of bond pairs	No. of lone pairs	Shape	Molecular geometry	Examples
	AB_3L	3	1	 pyramidal	 pyramidal	$NH_3, PF_3,$ $ClO_3^-, H_3O^+,$
	AB_2L_2	2	2	 Bent (V Shaped)	 Bent (V Shaped)	$H_2O, OF_2,$ SCl_2
5	AB_5	5	-	 Trigonal bipyramidal	 Trigonal bipyramidal	$PCl_5, ASF_5,$ SOF_4
	AB_4L	4	1	 Seesaw	 Seesaw	$SF_4, XeO_2F_2,$ $IF_4^+, IO_2F_2^-$
	AB_3L_2	3	2	 T shaped	 T Shaped	$BrF_3, ClF_3,$

Number of electron Pairs	Molecule	No. of bond pairs	No. of lone pairs	Shape	Molecular geometry	Examples
	AB_2L_3	2	3	 <p>Linear</p>	 <p>Linear</p>	XeF_2, I_3^-, IF_2^-
	AB_6	6	-	 <p>Octahedral</p>	 <p>Octahedral</p>	SF_6, IOF_5
6	AB_5L	5	1	 <p>Square pyramidal</p>	 <p>Square pyramidal</p>	$BrF_5, TeF_5^-, XeOF_4$
	AB_4L_2	4	2	 <p>Square planar</p>	 <p>Square Planar</p>	XeF_4, ICl_4^-
7	AB_7	7	-		 <p>pentagonal bi-pyramidal</p>	IF_7

Evaluate Yourself

6) Arrange the following in the decreasing order of Bond angle

- i) $\text{CH}_4, \text{H}_2\text{O}, \text{NH}_3$ ii) $\text{C}_2\text{H}_2, \text{BF}_3, \text{CCl}_4$

10.7 Valence Bond Theory

Heitler and London gave a theoretical treatment to explain the formation of covalent bond in hydrogen molecule on the basis of wave mechanics of electrons. It was further developed by Pauling and Slater. The wave mechanical treatment of VB theory is beyond the scope of this textbook. A simple qualitative treatment of VB theory for the formation of hydrogen molecule is discussed below.

Consider a situation wherein two hydrogen atoms (H_a and H_b) are separated by infinite distance. At this stage there is no interaction between these two atoms and the potential energy of this system is arbitrarily taken as zero. As these two atoms approach each other, in addition to the electrostatic attractive force between the nucleus and its own electron (purple arrows), the following new forces begins to operate.




Fig 10. 17 (a) VB theory for the formation of hydrogen molecule

The new attractive forces (green arrows) arise between

- nucleus of H_a and valence electron of H_b
- nucleus of H_b and the valence electron of H_a .

The new repulsive forces (red arrows) arise between

- the nucleus of H_a and H_b
- valence electrons of H_a and H_b .

The attractive forces tend to bring H_a and H_b together whereas the repulsive forces tends to push them apart. At the initial stage, as the two hydrogen atoms approach each other, the attractive forces are stronger than the repulsive forces and the potential energy decreases. A stage is reached where the net attractive forces are exactly balanced by repulsive forces and the potential energy of the system acquires a minimum energy.




Fig 10. 17 (b) VB theory for the formation of hydrogen molecule

At this stage, there is a maximum overlap between the atomic orbitals of H_a and H_b , and the atoms H_a and H_b are now said to be bonded together by a covalent bond. The internuclear distance at this stage gives the H-H bond length and is equal to 74 pm. The liberated energy is 436 kJ mol^{-1} and is known as bond energy. Since the energy is released during the bond formation, the resultant molecule is more stable. If the distance between the two atoms is decreased further, the repulsive forces dominate the attractive forces and the potential energy of the system sharply increases

10.7.1 Salient features of VB Theory:

- (i) When half filled orbitals of two atoms overlap, a covalent bond will be formed between them.
- (ii) The resultant overlapping orbital is occupied by the two electrons with opposite spins. For example, when H_2 is formed, the two 1s electrons of two hydrogen atoms get paired up and occupy the overlapped orbital.
- (iii) The strength of a covalent bond depends upon the extent of overlap of atomic orbitals. Greater the overlap, larger is the energy released and stronger will be the bond formed.

(iv) Each atomic orbital has a specific direction (except s-orbital which is spherical) and hence orbital overlap takes place in the direction that maximizes overlap

Let us explain the covalent bond formation in hydrogen, fluorine and hydrogen fluoride using VB theory.

10.8 Orbital Overlap

When atoms combine to form a covalent molecule, the atomic orbitals of the combining atoms overlap to form a covalent bond. The bond pair of electrons will occupy the overlapped region of the orbitals. Depending upon the nature of overlap we can classify the covalent bonding between the two atoms as sigma (σ) and pi (π) bonds.

10.8.1 Sigma and Pi bonds

When two atomic orbitals overlap linearly along the axis, the resultant bond is called a sigma (σ) bond. This overlap is also called 'head-on overlap' or 'axial overlap'. Overlap involving an s orbital (s-s and s-p overlaps) will always result in a sigma bond as the s orbital is spherical. Overlap between two p orbitals along the molecular axis will also result in sigma bond formation. When we consider x-axis as molecular axis, the p_x - p_x overlap will result in σ -bond.

When two atomic orbitals overlap sideways, the resultant covalent bond is called a pi (π) bond. When we consider x-axis as molecular axis, the p_y - p_y and p_z - p_z overlaps will result in the formation of a π -bond.

Following examples will be useful to understand the overlap:

10.8.2 Formation of hydrogen (H_2) Molecule

Electronic configuration of hydrogen atom is $1s^1$

During the formation of H_2 molecule, the $1s$ orbitals of two hydrogen atoms containing one unpaired electron with opposite spin overlap with each other along the internuclear axis. This overlap is called s-s overlap. Such axial overlap results in the formation of a σ -covalent bond.




Fig 10. 18 Formation of hydrogen molecule

Formation of fluorine molecule (F₂):

Valence shell electronic configuration of fluorine atom : $2s^2 2p_x^2, 2p_y^2, 2p_z^1$

When the half filled p_z orbitals of two fluorine overlaps along the z-axis, a σ -covalent bond is formed between them.




Fig 10.19 Formation of F₂ Molecule

Formation of HF molecule:

Electronic configuration of hydrogen atom is $1s^1$

Valence shell electronic configuration of fluorine atom : $2s^2 2p_x^2, 2p_y^2, 2p_z^1$

When half filled $1s$ orbital of hydrogen linearly overlaps with a half filled $2p_z$ orbital of fluorine, a σ -covalent bond is formed between hydrogen and fluorine.




Fig 10.20 Formation of HF Molecule

Formation of oxygen molecule (O₂):

Valence shell electronic configuration of oxygen atom : $2s^2 2p_x^2, 2p_y^1, 2p_z^1$

		π bond	σ bond
Oxygen 1	$1\downarrow$ $2s^2$	$1\downarrow$ $2p_x^2$	1 $2p_y^1$
Oxygen 2	$1\downarrow$ $2s^2$	$1\downarrow$ $2p_x^2$	1 $2p_z^1$

When the half filled p_z orbitals of two oxygen overlaps along the z-axis (considering molecular axis as z axis), a σ -covalent bond is formed between them. Other two half filled p_y orbitals of two oxygen atoms overlap laterally (sideways) to form a π -covalent bond between

the oxygen atoms. Thus, in oxygen molecule, two oxygen atoms are connected by two covalent bonds (double bond). The other two pair of electrons present in the $2s$ and $2p_x$ orbital do not involve in bonding and remains as lone pairs on the respective oxygen.




Fig 10. 21 Formation of π bond in O_2 Molecule

Evaluate Yourself

7) Bond angle in PH_4^+ is higher than in PH_3 why?

10.9 Hybridisation


Bonding in simple molecules such as hydrogen and fluorine can easily be explained on the basis of overlap of the respective atomic orbitals of the combining atoms. But the observed properties of polyatomic molecules such as methane, ammonia, beryllium chloride etc... cannot be explained on the basis of simple overlap of atomic orbitals. For example, it was experimentally proved that methane has a tetrahedral structure and the four C-H bonds are equivalent. This fact cannot be explained on the basis of overlap of atomic orbitals of hydrogen ($1s$) and the atomic orbitals of carbon with different energies ($2s^2 2p_x^2 2p_y 2p_z$).

In order to explain these observed facts, Linus Pauling proposed that the valence atomic orbitals in the molecules are different from those in isolated atom and he introduced the concept of hybridisation. Hybridisation is the process of mixing of atomic orbitals of the same atom with comparable energy to form equal number of new equivalent orbitals with same energy. The resultant orbitals are called hybridised orbitals and they possess maximum symmetry and definite orientation in space so as to minimize the force of repulsion between their electrons .

10.9.1 Types of hybridisation and geometry of molecules


sp Hybridisation:

Consider the bond formation in beryllium chloride. The ground state valence shell electronic configuration of Beryllium atom is $[\text{He}]2s^2 2p^0$



In BeCl_2 both the Be-Cl bonds are equivalent and it was observed that the molecule is linear. VB theory explain this observed behaviour by sp hybridisation. One of the paired electrons in the 2s orbital gets excited to 2p orbital and the electronic configuration at the excited state is shown.

Now, the 2s and 2p orbitals hybridise and produce two equivalent sp hybridised orbitals which have 50 % s-character and 50 % p-character. These sp hybridised orbitals are oriented in opposite direction as shown in the figure.



Overlap with orbital of chlorine

Each of the sp hybridized orbitals linearly overlap with $3p_z$ orbital of the chlorine to form a covalent bond between Be and Cl as shown in the Figure.





Fig 10.22 sp Hybridisation : BeCl_2


sp² Hybridisation:

Consider the bond formation in boron trifluoride. The ground state valence shell electronic configuration of Boron atom is [He]2s² 2p¹.



In the ground state boron has only one unpaired electron in the valence shell. In order to form three covalent bonds with fluorine atoms, three unpaired electrons are required. To achieve this, one of the paired electrons in the 2s orbital is promoted to the 2p_y orbital in the excited state.

In boron, the s orbital and two p orbitals (p_x and p_y) in the valence shell hybridises, to generate three equivalent sp² orbitals as shown in the Figure. These three orbitals lie in the same xy plane and the angle between any two orbitals is equal to 120°



Overlap with 2p_z orbitals of fluorine:

The three sp² hybridised orbitals of boron now overlap with the 2p_z orbitals of fluorine (3 atoms). This overlap takes place along the axis as shown below.





Fig 10.23 sp² Hybridation : BF₃

sp³ Hybridisation:

sp³ hybridisation can be explained by considering methane as an example. In methane molecule the central carbon atom bound to four hydrogen atoms. The ground state valence shell electronic configuration of carbon is [He]2s² 2p_x¹ 2p_y¹ 2p_z⁰.



In order to form four covalent bonds with the four hydrogen atoms, one of the paired electrons in the 2s orbital of carbon is promoted to its 2p_z orbital in the excited state. The one 2s orbital and the three 2p orbitals of carbon mix to give four equivalent sp³ hybridised orbitals. The angle between any two sp³ hybridised orbitals is 109° 28'

Overlap with 1s orbitals of hydrogen:

The 1s orbitals of the four hydrogen atoms overlap linearly with the four sp³ hybridised orbitals of carbon to form four C-H σ-bonds in the methane molecule, as shown below.






Fig 10.24 sp³ Hybridisation : CH₄

sp³d Hybridisation:

In the molecules such as PCl₅, the central atom phosphorus is covalently bound to five chlorine atoms. Here the atomic orbitals of phosphorous undergoes sp³d hybridisation which involves its one 3s orbital, three 3p orbitals and one vacant 3d orbital (d_{z²}). The ground state electronic configuration of phosphorous is [Ne]3s² 3p_x¹ 3p_y¹ 3p_z¹ as shown below.



One of the paired electrons in the 3s orbital of phosphorous is promoted to one of its vacant 3d orbital (d_{z²}) in the excited state. One 3s orbital, three 3p orbitals and one 3d_{z²} orbital of phosphorus atom mixes to give five equivalent sp³d hybridised orbitals. The orbital geometry of sp³d hybridised orbitals is trigonal bi-pyramidal as shown in the figure 10.25.


Overlap with 3p_z orbitals of chlorine:

The 3p_z orbitals of the five chlorine atoms linearly overlap along the axis with the five sp³d hybridised orbitals of phosphorous to form the five P-Cl σ-bonds, as shown below.

Fig 10.25 sp³d Hybridisation : PCl₅

Bonding in Ethylene:

The bonding in ethylene can be explained using hybridisation concept. The molecular formula of ethylene is C_2H_4 . The valency of carbon is 4. The electronic configuration of valence shell of carbon in ground state is $[He]2s^2 2p_x^1 2p_y^1 2p_z^0$. To satisfy the valency of carbon promote an electron from $2s$ orbital to $2p_z$ orbital in the excited state.



In ethylene both the carbon atoms undergoes sp^2 hybridisation involving $2s$, $2p_x$ and $2p_y$ orbitals, resulting in three equivalent sp^2 hybridised orbitals lying in the xy plane at an angle of 120° to each other. The unhybridised $2p_z$ orbital lies perpendicular to the xy plane.

Formation of sigma bond:

One of the sp^2 hybridised orbitals of each carbon lying on the molecular axis (x -axis) linearly overlaps with each other resulting in the formation of a C-C sigma bond. Other two sp^2 hybridised orbitals of both carbons linearly overlap with the four $1s$ orbitals of four hydrogen atoms leading to the formation of two C-H sigma bonds on each carbon.

Formation of Pi (π) bond:

The unhybridised $2p_z$ orbital of both carbon atoms can overlap only sideways as they are not in the molecular axis. This lateral overlap results in the formation of a pi (π) bond between the two carbon atoms as shown in the figure.




Fig 10.27 sp^2 Hybridisation : C_2H_4

Bonding in acetylene:

Similar to ethylene, the bonding in acetylene can also be explained using hybridisation concept. The molecular formula of acetylene is C_2H_2 . The electronic configuration of valence shell of carbon in ground state is $[He]2s^2 2p_x^1 2p_y^1 2p_z^0$. To satisfy the valency of carbon promote an electron from 2s orbital to $2p_z$ orbital in the excited state.

In acetylene molecule, both the carbon atoms are in sp hybridised state. The $2s$ and $2p_x$ orbitals, resulting in two equivalent sp hybridised orbitals lying in a straight line along the molecular axis (x-axis). The unhybridised $2p_y$ and $2p_z$ orbitals lie perpendicular to the molecular axis.

Formation of sigma bond:

One of the two sp hybridised orbitals of each carbon linearly overlaps with each other resulting in the formation a C-C sigma bond. The other sp hybridised orbital of both carbons linearly overlap with the two $1s$ orbitals of two hydrogen atoms leading to the formation of one C-H sigma bonds on each carbon.

Formation of pi bond:

The unhybridised $2p_y$ and $2p_z$ orbitals of each carbon overlap sideways. This lateral overlap results in the formation of two pi bonds (p_y-p_y and p_z-p_z) between the two carbon atoms as shown in the figure.




Fig 10.28 sp Hybridisation in acetylene: C_2H_2

Evaluate Yourself

8) Explain the bond formation in SF_4 and CCl_4 using hybridisation concept.

Evaluate Yourself

9) The observed bond length of N_2^+ is larger than N_2 while the bond length in NO^+ is less than in NO . Why?

10.10 Molecular orbital theory

Lewis concept and valence bond theory qualitatively explains the chemical bonding and molecular structure. Both approaches are inadequate to describe some of the observed properties of molecules. For example, these theories predict that oxygen is diamagnetic. However, it was observed that oxygen in liquid form was attracted towards the poles of strong magnet, indicating that oxygen is paramagnetic. As both these theories treated the bond formation in terms of electron pairs and hence they fail to explain the bonding nature of paramagnetic molecules. F. Hund and Robert. S. Mulliken developed a bonding theory called molecular orbital theory which explains the magnetic behaviour of molecules.

The salient features of Molecular orbital Theory (MOT):

- When atoms combine to form molecules, their individual atomic orbitals lose their identity and form new orbitals called molecular orbitals.
- The shapes of molecular orbitals depend upon the shapes of combining atomic orbitals.
- The number of molecular orbitals formed is the same as the number of combining atomic orbitals. Half the number of molecular orbitals formed will have lower energy than the corresponding atomic orbital, while the remaining molecular orbitals will have higher energy. The molecular orbital with lower energy is called bonding molecular orbital and the one with higher energy is called anti-bonding molecular orbital. The bonding molecular orbitals are represented as σ (Sigma), π (pi), δ (delta) and the corresponding antibonding orbitals are denoted as σ^* , π^* and δ^* .
- The electrons in a molecule are accommodated in the newly formed molecular orbitals. The filling of electrons in these orbitals follows Aufbau's principle, Pauli's exclusion principle and Hund's rule as in the case of filling of electrons in atomic orbitals.
- Bond order gives the number of covalent bonds between the two combining atoms. The bond order of a molecule can be calculated using the following equation

$$\text{Bond order} = \frac{N_b - N_a}{2}$$

Where,

N_b = Total number of electrons present in the bonding molecular orbitals

N_a = Total number of electrons present in the antibonding molecular orbitals and

A bond order of zero value indicates that the molecule doesn't exist.

10.10.1 Linear combination of atomic orbitals

The wave functions for the molecular orbitals can be obtained by solving Schrödinger wave equation for the molecule. Since solving the Schrödinger equation is too complex, approximation methods are used to obtain the wave function for molecular orbitals. The most common method is the linear combination of atomic orbitals (LCAO).


We know that the atomic orbitals are represented by the wave function Ψ . Let us consider two atomic orbitals represented by the wave function ψ_A and ψ_B with comparable energy, combines to form two molecular orbitals. One is bonding molecular orbital (ψ_{bonding}) and the other is antibonding molecular orbital ($\psi_{\text{antibonding}}$). The wave functions for these two molecular orbitals can be obtained by the linear combination of the atomic orbitals ψ_A and ψ_B as below.

$$\Psi_{\text{bonding}} = \psi_A + \psi_B$$

$$\Psi_{\text{antibonding}} = \psi_A - \psi_B$$

The formation of bonding molecular orbital can be considered as the result of constructive interference of the atomic orbitals and the formation of anti-bonding molecular orbital can be the result of the destructive interference of the atomic orbitals. The formation of the two molecular orbitals from two 1s orbitals is shown below.

Constructive interaction: The two 1s orbitals are in phase and have the same sign.



Destructive interaction The two 1s Orbitals are out of phase




Fig 10.29 Linear Combination of atomic orbitals

10.10.2 Bonding in some Homonuclear di-atomic molecules:




Fig 10.30 MO Diagram for H₂ molecule

Molecular orbital diagram of hydrogen molecule (H₂)

Electronic configuration of H atom is $1s^1$
 Electronic configuration of H₂ molecule σ_{1s}^2

$$\text{Bond order} = \frac{N_b - N_a}{2} = \frac{2 - 0}{2} = 1$$

Molecule has *no unpaired electrons*.
 Hence, it is **diamagnetic**.




Fig 10.31 MO Diagram for Li₂ molecule

Molecular orbital diagram of lithium molecule (Li₂)

Electronic configuration of Li atom is $1s^2 2s^1$

Electronic configuration of Li₂ molecule $\sigma_{1s}^2, \sigma_{1s}^{*2}, \sigma_{2s}^2$

$$\text{Bond order} = \frac{N_b - N_a}{2} = \frac{4 - 2}{2} = 1$$

Molecule has *no unpaired electrons*.
 Hence it is **diamagnetic**.




Fig 10.32 MO Diagram for B₂ molecule

Molecular orbital diagram of boron molecule (B₂)

Electronic configuration of B atom is $1s^2 2s^2 2p^1$

Electronic configuration of B₂ molecule $\sigma_{1s}^2, \sigma_{1s}^{*2}, \sigma_{2s}^2, \sigma_{2s}^{*2}, \pi_{2p_y}^1, \pi_{2p_z}^1$

$$\text{Bond order} = \frac{N_b - N_a}{2} = \frac{6 - 4}{2} = 1$$

Molecule has *two unpaired electrons*.
 Hence it is **paramagnetic**.






Fig 10.33 MO Diagram for C_2 molecule

Molecular orbital diagram of carbon molecule (C_2)

Electronic configuration of C atom is



Electronic configuration of C_2 molecule



$$\text{Bond order} = \frac{N_b - N_a}{2} = \frac{8 - 4}{2} = 2$$

Molecule has *no unpaired electrons*. Hence, it is **diamagnetic**.






Fig 10.34 MO Diagram for N_2 molecule

Molecular orbital diagram of nitrogen molecule (N_2)

Electronic configuration of N atom is



Electronic configuration of N_2 molecule



$$\text{Bond order} = \frac{N_b - N_a}{2} = \frac{10 - 4}{2} = 3$$

Molecule has *no unpaired electrons*. Hence, it is **diamagnetic**.






Fig 10.35 MO Diagram for O_2 molecule

Molecular orbital diagram of oxygen molecule (O_2)

Electronic configuration of O atom is



Electronic configuration of O_2 molecule



$$\text{Bond order} = \frac{N_b - N_a}{2} = \frac{10 - 6}{2} = 2$$

Molecule has *two unpaired electrons*. Hence, it is **paramagnetic**.

10.10.3 Bonding in some Heteronuclear di-atomic molecules




Fig 10.36 MO Diagram for CO molecule

Molecular orbital diagram of Carbon monoxide molecule (CO)

Electronic configuration of C atom is $1s^2 2s^2 2p^2$

Electronic configuration of O atom is $1s^2 2s^2 2p^4$

Electronic configuration of CO molecule $\sigma_{1s}^2, \sigma_{1s}^{*2}, \sigma_{2s}^2, \sigma_{2s}^{*2}, \pi_{2p_y}^2, \pi_{2p_z}^2, \sigma_{2p_x}^2$

$$\text{Bond order} = \frac{N_b - N_a}{2} = \frac{10 - 4}{2} = 3$$

Molecule has *no unpaired electrons*. Hence, it is **diamagnetic**.




Fig 10.36 MO Diagram for NO molecule

Molecular orbital diagram of Nitric oxide molecule (NO)

Electronic configuration of N atom is $1s^2 2s^2 2p^3$

Electronic configuration of O atom is $1s^2 2s^2 2p^4$

Electronic configuration of NO molecule $\sigma_{1s}^2, \sigma_{1s}^{*2}, \sigma_{2s}^2, \sigma_{2s}^{*2}, \pi_{2p_y}^2, \pi_{2p_z}^2, \sigma_{2p_x}^2, \pi_{2p_y}^{*1}$

$$\text{Bond order} = \frac{N_b - N_a}{2} = \frac{10 - 5}{2} = 2.5$$

Molecule has *one unpaired electron*. Hence, it is **paramagnetic**.

Evaluate Yourself

10) Draw the MO diagram for acetylide ion C_2^{2-} and calculate its bond order.

Metallic bonding

Metals have some special properties of lustre, high density, high electrical and thermal conductivity, malleability and ductility, and high melting and boiling points. The forces that keep the atoms of the metal so closely in a metallic crystal constitute what is generally known as the metallic bond. The metallic bond is not just an electrovalent bond(ionic bond), as the latter is formed between atoms of different electro negativities. Similarly, the metallic bond is not a covalent bond,as the metal atoms do not have sufficient number of valence electrons for mutual sharing with 8 or 12 neighboring metal atoms in a crystal. So, we have to search for a new theory to explain metallic bond. The first successful theory is due to Drude and Lorentz, which regards metallic crystal as an assemblage of positive ions immersed in a gas of free electrons. The free electrons are due to ionization of the valence electrons of the atoms of the metal. As the valence electrons of the atoms are freely shared by all the ions in the crystal, the metallic bonding is also referred to as electronic bonding. As the free electrons repel each other, they are uniformly distributed around the metal ions. Many physical properties of the metals can be explained by this theory, nevertheless there are exceptions.

The electrostatic attraction between the metal ions and the free electrons yields a three-dimensional close packed crystal with a large number of nearest metal ions. So, metals have high density. As the close packed structure contains many slip planes along which movement can occur during mechanical loading, the metal acquires

ductility. Pure metals can undergo 40 to 60% elongation prior to rupturing under mechanical loading. As each metal ion is surrounded by electron cloud in all directions, the metallic bonding has no directional properties.

As the electrons are free to move around the positive ions, the metals exhibit high electrical and thermal conductivity. The metallic luster is due to reflection of light by the electron cloud. As the metallic bond is strong enough, the metal atoms are reluctant to break apart into a liquid or gas, so the metals have high melting and boiling points.

The bonding in metal is better treated by Molecular orbital theory. As per this theory, the atomic orbitals of large number of atoms in a crystal overlap to form numerous bonding and antibonding molecular orbitals without any band gap. The bonding molecular orbitals are completely filled with an electron pair in each, and the antibonding molecular orbitals are empty. Absence of band gap accounts for high electrical conductivity of metals. High thermal conductivity is due to thermal excitation of many electrons from the valence band to the conductance band. With an increase in temperature, the electrical conductivity decreases due to vigorous thermal motion of lattice ions that disrupts the uniform lattice structure, that is required for free motion of electrons within the crystal. Most metals are black except copper, silver and gold. It is due to absorption of light of all wavelengths. Absorption of light of all wavelengths is due to absence of bandgap in metals.



SUMMARY

In molecules, atoms are held together by attractive forces, called chemical bonds. Kossel and Lewis are the first people to provide a logical explanation for chemical bonding. They proposed that atoms try to attain the nearest noble gas electronic configuration by losing, gaining or sharing one or more electrons during the bond formation. The noble gases contain eight electrons in their valance shell which is considered to be stable electronic configuration. The idea of Kossel – Lewis approach to chemical bond lead to the octet rule, which states that “the atoms transfer or share electrons so that all atoms involved in chemical bonding obtain 8 electrons in their outer shell (valance shell)”.

There are different types of chemical bonds. In compounds such as sodium chloride, the sodium atom loses an electron which is accepted by the chlorine atom resulting in the formation of Na^+ and Cl^- ions. These two ions are held together by the electrostatic attractive forces. This type of chemical bond is known as ionic bonds or electrovalent bonds. In certain compounds, instead of the complete transfer of electrons, the electrons are shared by both the bonding atoms. The two combining atoms are held together by their mutual attraction towards the shared electrons. This type of bond is called covalent bonding. In addition, there also another bond type known as coordinate covalent bonds, where the shared electrons of a covalent bond are provided by only one of the combining atoms. Metallic bonding is another type of bonding which is observed in metals.

Lewis theory in combination with VSEPR theory will be useful in predicting the shape of molecules. According to this theory, the shape of the molecules depends on the number of valance shell electron pair (lone pairs and bond pairs) around the central atom. Each pair of valance electrons around the central atom repels each other and hence, they are located as far away as possible in three-dimensional space to minimise the repulsion between them.

Heitler and London gave a theoretical treatment to explain the formation of covalent bond in hydrogen molecule on the basis of wave mechanics of electrons. It was further developed by Pauling and Slater. According to this theory when half-filled orbitals of two atoms overlap, a covalent bond will be formed between them. Linus Pauling introduced the concept of hybridisation. Hybridisation is the process of mixing of atomic orbitals of the same atom with comparable energy to form equal number of new equivalent orbitals with same energy. There are different types of hybridization such as sp , sp^2 , sp^3 , sp^3d^2 etc..

F. Hund and Robert. S. Mulliken developed a bonding theory called molecular orbital theory. According to this theory, when atoms combines to form molecules, their individual atomic orbitals lose their identity and forms new orbitals called molecular orbitals. The filling of electrons in these orbitals follows Aufbau's principle, Pauli's exclusion principle and Hund's rule as in the case of filling of electrons in atomic orbitals.

Evaluation







I. Choose the best answer.

- In which of the following Compounds does the central atom obey the octet rule?
a) XeF_4 b) AlCl_3
c) SF_6 d) SCl_2
- In the molecule $\text{O}_A = \text{C} = \text{O}_B$, the formal charge on O_A , C and O_B are respectively.
a) -1, 0, +1 b) +1, 0, -1
c) -2, 0, +2 d) 0, 0, 0
- Which of the following is electron deficient?
a) PH_3 b) $(\text{CH}_3)_2$
c) BH_3 d) NH_3
- Which of the following molecule contain no π bond?
a) SO_2 b) NO_2
c) CO_2 d) H_2O
- The ratio of number of sigma (σ) and pi (π) bonds in 2-butyne is
a) 8/3 b) 5/3
c) 8/2 d) 9/2
- Which one of the following is the likely bond angles of sulphur tetrafluoride molecule?
a) $120^\circ, 80^\circ$ b) $109^\circ, 28^\circ$
c) 90° d) $89^\circ, 117^\circ$
- Assertion:** Oxygen molecule is paramagnetic.
Reason : It has two unpaired electron in its bonding molecular orbital
a) both assertion and reason are true and reason is the correct explanation of assertion
b) both assertion and reason are true but reason is not the correct explanation of assertion
c) assertion is true but reason is false
d) Both assertion and reason are false
- According to Valence bond theory, a bond between two atoms is formed when
a) fully filled atomic orbitals overlap
b) half filled atomic orbitals overlap
c) non-bonding atomic orbitals overlap
d) empty atomic orbitals overlap
- In ClF_3 , NF_3 and BF_3 molecules the chlorine, nitrogen and boron atoms are
a) sp^3 hybridised
b) sp^3, sp^3 and sp^2 respectively
c) sp^2 hybridised
d) sp^3d , sp^3 and sp^2 hybridised respectively
- When one s and three p orbitals hybridise,
a) four equivalent orbitals at 90° to each other will be formed
b) four equivalent orbitals at $109^\circ 28'$ to each other will be formed.





c) four equivalent orbitals, that are lying the same plane will be formed

d) none of these




11. Which of these represents the correct order of their increasing bond order.




12. Hybridisation of central atom in PCl_5 involves the mixing of orbitals.



13. The correct order of O-O bond length in hydrogen peroxide, ozone and oxygen is



14. Which one of the following is diamagnetic?



15. Bond order of a species is 2.5 and the number of electrons in its bonding molecular orbital is found to be 8. The no. of electrons in its antibonding molecular orbital is

a) three b) four

c) Zero d) can not be calculated from the given information.

16. Shape and hybridisation of IF_5 are

a) Trigonal bipyramidal, sp^3d^2

b) Trigonal bipyramidal, sp^3d

c) Square pyramidal, sp^3d^2

d) Octahedral, sp^3d^2

17. Pick out the incorrect statement from the following


a) sp^3 hybrid orbitals are equivalent and are at an angle of $109^\circ 28'$ with each other

b) dsp^2 hybrid orbitals are equivalent and bond angle between any two of them is 90°




c) All five sp^3d hybrid orbitals are not equivalent out of these five sp^3d hybrid orbitals, three are at an angle of 120° , remain two are perpendicular to the plane containing the other three

d) none of these

18. The molecules having same hybridisation, shape and number of lone pairs of electrons are



19. In which of the following molecules / ions BF_3 , NO_2^- , H_2O the central atom is sp^2 hybridised?



20. Some of the following properties of two species, NO_3^- and H_3O^+ are described below. which one of them is correct?
- dissimilar in hybridisation for the central atom with different structure.
 - isostructural with same hybridisation for the Central atom.
 - different hybridisation for the central atom with same structure
 - none of these
21. The types of hybridisation on the five carbon atom from right to left in the, 2,3 pentadiene.
- $\text{sp}^3, \text{sp}^2, \text{sp}, \text{sp}^2, \text{sp}^3$
 - $\text{sp}^3, \text{sp}, \text{sp}, \text{sp}, \text{sp}^3$
 - $\text{sp}^2, \text{sp}, \text{sp}^2, \text{sp}^2, \text{sp}^3$
 - $\text{sp}^3, \text{sp}^3, \text{sp}^2, \text{sp}^3, \text{sp}^3$
22. Xe F_2 is isostructural with
- SbCl_2
 - BaCl_2
 - TeF_2
 - ICl_2^-
23. The percentage of s-character of the hybrid orbitals in methane, ethane, ethene and ethyne are respectively
- 25, 25, 33.3, 50
 - 50, 50, 33.3, 25
 - 50, 25, 33.3, 50
 - 50, 25, 25, 50
24. Of the following molecules, which have shape similar to carbon dioxide?
- SnCl_2
 - NO_2
 - C_2H_2
 - All of these.
25. According to VSEPR theory, the repulsion between different parts of electrons obey the order.
- $\text{l.p} - \text{l.p} > \text{b.p} - \text{b.p} > \text{l.p} - \text{b.p}$
 - $\text{b.p} - \text{b.p} > \text{b.p} - \text{l.p} > \text{l.p} - \text{b.p}$
 - $\text{l.p} - \text{l.p} > \text{b.p} - \text{l.p} > \text{b.p} - \text{b.p}$
 - $\text{b.p} - \text{b.p} > \text{l.p} - \text{l.p} > \text{b.p} - \text{l.p}$
26. Shape of ClF_3 is
- Planar triangular
 - Pyramidal
 - 'T' Shaped
 - none of these
27. Non- Zero dipole moment is shown by
- CO_2
 - p-dichlorobenzene
 - carbontetrachloride
 - water.
28. Which of the following conditions is not correct for resonating structures?
- the contributing structure must have the same number of unpaired electrons
 - the contributing structures should have similar energies
 - the resonance hybrid should have higher energy than any of the contributing structure.
 - none of these
29. Among the following, the compound that contains, ionic, covalent and Coordinate linkage is
- NH_4Cl
 - NH_3

c) NaCl d) none of these

30. CaO and NaCl have the same crystal structure and approximately the same radii. If U is the lattice energy of NaCl, the approximate lattice energy of CaO is

a) U b) $2U$

c) $U/2$ d) $4U$

II. Write brief answer to the following questions.

31. Define the following

i) Bond order

ii) Hybridisation

iii) σ - bond

32. What is a pi bond?

33. In CH_4 , NH_3 and H_2O , the central atom undergoes sp^3 hybridisation - yet their bond angles are different. why?

34. Explain sp^2 hybridisation in BF_3

35. Draw the M.O diagram for oxygen molecule calculate its bond order and show that O_2 is paramagnetic.

36. Draw MO diagram of CO and calculate its bond order.

37. What do you understand by Linear combination of atomic orbitals in MO theory.

38. Discuss the formation of N_2 molecule using MO Theory

39. What is dipole moment?

40. Linear form of carbondioxide molecule has two polar bonds. yet the molecule has Zero dipole moment why?

41. Draw the Lewis structures for the following species.

i) NO_3^- ii) SO_4^{2-}

iii) HNO_3 iv) O_3

42. Explain the bond formation in BeCl_2 and MgCl_2 .

43. Which bond is stronger σ or π ? Why?

44. Define bond energy.

45. Hydrogen gas is diatomic where as inert gases are monoatomic - explain on the basis of MO theory.

46. What is Polar Covalent bond? explain with example.

47. Considering x- axis as molecular axis, which out of the following will form a sigma bond.

i) $1s$ and $2p_y$ ii) $2p_x$ and $2p_x$

iii) $2p_x$ and $2p_z$ iv) $1s$ and $2p_z$

48. Explain resonance with reference to carbonate ion?

49. Explain the bond formation in ethylene and acetylene.

50. What type of hybridisations are possible in the following geometries?

a) octahedral

b) tetrahedral

c) square planer.

51. Explain VSEPR theory. Applying this theory to predict the shapes of IF_7 and SF_6

52. CO_2 and H_2O both are triatomic molecule but their dipole moment values are different. Why?

53. Which one of the following has highest bond order?


N_2 , N_2^+ or N_2^-

54. Explain the covalent character in ionic bond.

55. Describe fajan's rule.



CONCEPT MAP





Shapes of Molecule (VSEPR Theory)


By using this tool, you will be able to create and visualise molecules with different number of bonds and lone pairs.

Please go to the URL
https://phet.colorado.edu/sims/html/molecule-shapes/latest/molecule-shapes_en.html
(or)
Scan the QR code on the right side



Steps

- Open the Browser and type the URL given (or) Scan the QR Code.
- Double click the 'models' icon given in this page. Now you will see the web page as shown in the figure.
- Now, by selecting the number of bond pairs and lone pairs using the options in the box 3 and 4 effectively, you visualize the corresponding molecular geometry.
- You can see the angles between the bonds selecting the "Show bond angles" indicated in box 5.
- You can also visualize the molecular geometry and the electron geometry by selecting corresponding option in box 2.



Structure of simple molecules:

By selecting the real molecules icon in indicated in the box 6, you will be able to see a screen as shown below. In this screen, use the dropdown menu indicated in box 7 you can select a molecule. Use the option in the box 8 you can visualise the lone pairs and bond pairs in the molecules.