

# Saitechinfo NEET-JEE Academy



## Definite Integral as the Limit of a Sum

The definite integral is a fundamental concept in calculus that connects the discrete summation of areas under a curve with continuous accumulation. Here's a structured lecture note on this concept:

---

### 1. Introduction to Definite Integrals

The definite integral of a function  $f(x)$  over an interval  $[a, b]$  represents the net area under the curve  $y = f(x)$  from  $x = a$  to  $x = b$ .

$$\int_a^b f(x) dx$$

This integral can be interpreted as the limit of a sum of areas of rectangles as the number of rectangles approaches infinity.

---

### 2. Approximating Area Using Riemann Sums

To approximate the area under the curve:

1. Divide the interval  $[a, b]$  into  $n$  subintervals of equal width:

$$\Delta x = \frac{b - a}{n}$$

2. In each subinterval, choose a point  $x_i^*$  (e.g., left endpoint, right endpoint, or midpoint).
3. Compute the sum of the areas of rectangles formed by  $f(x_i^*)$  (height) and  $\Delta x$  (width):

$$S_n = \sum_{i=1}^n f(x_i^*) \Delta x$$

---

### 3. Limit of Riemann Sum

As  $n \rightarrow \infty$ , the width of the rectangles ( $\Delta x$ ) approaches zero, and the sum converges to the exact area under the curve:

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x$$

This is the definition of the definite integral as a limit of a Riemann sum.

---

### 4. Examples

**Example 1: Compute the Definite Integral**

Find  $\int_0^1 x^2 dx$  using the definition of a definite integral.

1. Divide  $[0, 1]$  into  $n$  subintervals:

$$\Delta x = \frac{1 - 0}{n} = \frac{1}{n}$$

2. Use right endpoints:  $x_i^* = \frac{i}{n}$  for  $i = 1, 2, \dots, n$ .

3. The Riemann sum becomes:

$$S_n = \sum_{i=1}^n f\left(\frac{i}{n}\right) \Delta x = \sum_{i=1}^n \left(\frac{i}{n}\right)^2 \cdot \frac{1}{n}$$

4. Simplify:

$$S_n = \frac{1}{n^3} \sum_{i=1}^n i^2$$

5. Use the formula for the sum of squares:

$$\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$$

6. Substitute and simplify:

$$S_n = \frac{1}{n^3} \cdot \frac{n(n+1)(2n+1)}{6} = \frac{(n+1)(2n+1)}{6n^2}$$

7. Take the limit as  $n \rightarrow \infty$ :

$$\int_0^1 x^2 dx = \lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \frac{(n+1)(2n+1)}{6n^2} = \frac{1}{3}$$

**Example 2: Verify  $\int_1^2 3x^2 dx$** 

1. Divide  $[1, 2]$  into  $n$  subintervals:

$$\Delta x = \frac{2 - 1}{n} = \frac{1}{n}$$

2. Use left endpoints:  $x_i^* = 1 + \frac{i-1}{n}$  for  $i = 1, 2, \dots, n$ .

3. The Riemann sum becomes:

$$S_n = \sum_{i=1}^n f(x_i^*) \Delta x = \sum_{i=1}^n 3 \left(1 + \frac{i-1}{n}\right)^2 \cdot \frac{1}{n}$$

4. Expand  $\left(1 + \frac{i-1}{n}\right)^2$ , simplify the sum, and take the limit as  $n \rightarrow \infty$ .

Result:

$$\int_1^2 3x^2 dx = [x^3]_1^2 = 2^3 - 1^3 = 8 - 1 = 7$$

---

## 5. Key Points

1. The definite integral calculates the exact area under a curve by taking the limit of Riemann sums.
2. Riemann sums depend on the choice of  $x_i^*$ , but all approaches lead to the same integral value as  $n \rightarrow \infty$ .

These examples illustrate the concept and method of evaluating definite integrals using the limit of a sum.